

# WEST Search History





DATE: Saturday, September 17, 2005

<u>Hide?</u>	<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>
	<i>DB=DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L28	L27 and l26	62
<input type="checkbox"/>	L27	toxin	7426
<input type="checkbox"/>	L26	quinguestriatus or leiurus or scorpion	652
	<i>DB=JPAB; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L25	quinguestriatus or leiurus or scorpion	6
	<i>DB=EPAB; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L24	L23 and l22	9
<input type="checkbox"/>	L23	toxin	1177
<input type="checkbox"/>	L22	quinguestriatus or leiurus or scorpion	29
	<i>DB=USOC; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L21	L20 same l19	0
<input type="checkbox"/>	L20	toxin	463
<input type="checkbox"/>	L19	quinguestriatus or leiurus or scorpion	28
<input type="checkbox"/>	L18	potassium or k+	105038
	<i>DB=PGPB; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L17	L15 and (potassium or k+)	109
<input type="checkbox"/>	L15	(L14 or l12) same l13	171
<input type="checkbox"/>	L14	scorpion	835
<input type="checkbox"/>	L13	toxin	22638
<input type="checkbox"/>	L12	quinguestriatus or leiurus	43
	<i>DB=USPT; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L11	l7 and l6 not l9	61
<input type="checkbox"/>	L10	l7 and l3 not l8	33
<input type="checkbox"/>	L9	l7 same l6	34
<input type="checkbox"/>	L8	L7 same l3	3
<input type="checkbox"/>	L7	potassium or k+	282322
<input type="checkbox"/>	L6	L5 not l3	208
<input type="checkbox"/>	L5	l4 same l2	258
<input type="checkbox"/>	L4	scorpion	1172
<input type="checkbox"/>	L3	l1 same L2	53
<input type="checkbox"/>	L2	toxin	22671

☐ L1 quinquestriatus or leiurus

79

END OF SEARCH HISTORY

## SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1951-2005/Sep 16 (c) format only 2005 Dialog

File 5:Biostis Previews(R) 1969-2005/Sep W2 (c) 2005 BIOSIS

File 357:Derwent Biotech Res. 1982-2005/Sep W3 (c) 2005 Thomson Derwent &amp; ISI

Set Items Description  
 S1 768 QUINQUESTRIATUS  
 S2 231300 TOXIN  
 S3 336 S1 AND S2  
 S4 3963 CHARYBDOTOXIN  
 S5 62 S3 AND S4  
 S6 62 ID (sorted in duplicate order)  
 S7 1790727 HOMOLOG? OR ANALOG? OR FAMILY  
 S8 18 S6 AND S7  
 S9 18 ID (sorted in duplicate order)  
 S10 838 LEIURUS  
 S11 119 S10 NOT S1  
 S12 1 S11 AND S2 AND S4 AND S7  
 S13 2 S11 AND S2 AND S4  
 S14 526 (CYSTINE OR CYSTEINE)(W)KNOT  
 S15 0 S1 AND S14  
 S16 2 S4 AND S14  
 S17 7240 SCORPION  
 S18 81 S17 AND S2 AND S4 AND S7 NOT S8  
 S19 81 ID (sorted in duplicate order)

S24 29 ID (sorted in duplicate order)  
 S25 9 S1 AND S22  
 S26 9 ID (sorted in duplicate order)

Ref Items Index-term  
 E1 1 AU=PRESMANES MORALES S  
 E2 2 AU=PRESME N  
 E3 1 \*AU=PRESNAIL J  
 E4 19 AU=PRESNAIL J K  
 E5 10 AU=PRESNAIL JAMES K  
 E6 1 AU=PRESNAIL JIM  
 E7 1 AU=PRESNAJDER P  
 E8 2 AU=PRESNAL B  
 E9 3 AU=PRESNAL B P  
 E10 1 AU=PRESNAL J K  
 E11 1 AU=PRESNALL C C  
 E12 2 AU=PRESNALL D

S27 30 E3-E5  
 S28 1 S1 AND S27  
 S29 3 S17 AND S27

Ref Items Index-term  
 E1 1 AU=HERRMANN PHIL U  
 E2 6 AU=HERRMANN PIA  
 E3 850 \*AU=HERRMANN R  
 E4 15 AU=HERRMANN R A  
 E5 4 AU=HERRMANN R B  
 E6 3 AU=HERRMANN R E  
 E7 1 AU=HERRMANN R F  
 E8 309 AU=HERRMANN R G  
 E9 5 AU=HERRMANN R J  
 E10 7 AU=HERRMANN R K  
 E11 39 AU=HERRMANN R L  
 E12 11 AU=HERRMANN R M  
 E13 4 AU=HERRMANN R O  
 E14 91 AU=HERRMANN R P  
 E15 5 AU=HERRMANN R R  
 E16 3 AU=HERRMANN R S  
 E17 6 AU=HERRMANN R W  
 E18 1 AU=HERRMANN R-M  
 E19 12 AU=HERRMANN RAFAEL  
 E20 2 AU=HERRMANN RAFI  
 E21 1 AU=HERRMANN RAINALD  
 E22 16 AU=HERRMANN RALF  
 E23 3 AU=HERRMANN RALPH  
 E24 4 AU=HERRMANN REBEKAH R

S20 1366 E3-E19  
 S21 1 AU=HERRMANN, R G'  
 S22 1367 S21 OR S20  
 S23 29 S17 AND S22

E7 1 AU=LU AIYUN  
 E8 1 AU=LU AL  
 E9 1 AU=LU AL-LING  
 E10 1 AU=LU ALAN Q  
 E11 2 AU=LU ALAN QING  
 E12 1 AU=LU ALAN QUIN  
 E13 23 AU=LU ALBERT  
 E14 2 AU=LU ALBERT L  
 E15 1 AU=LU ALEX  
 E16 4 AU=LU ALICE  
 E17 1 AU=LU ALLAN T K  
 E18 2 AU=LU ALLEN  
 E19 6 AU=LU ALLEN P

S31 25 E13,E14  
 S32 244 S30 OR S31  
 S33 2 S1 AND S32  
 S34 4 S17 AND S32

Ref Items Index-term  
 E1 1 AU=WONG IVY HN  
 E2 1 AU=WONG IVY S L  
 E3 1352 \*AU=WONG J  
 E4 25 AU=WONG J A  
 E5 2 AU=WONG J A L  
 E6 4 AU=WONG J A-L  
 E7 124 AU=WONG J B  
 E8 95 AU=WONG J C  
 E9 30 AU=WONG J C F  
 E10 9 AU=WONG J C H  
 E11 2 AU=WONG J C K  
 E12 5 AU=WONG J C L

S35 361 S3,AU=WONG J A'

Ref Items Index-term  
 E1 1 AU=WONG JAFFE  
 E2 3 AU=WONG JAIME A  
 E3 41 \*AU=WONG JAMES  
 E4 1 AU=WONG JAMES B  
 E5 1 AU=WONG JAMES CHING  
 E6 6 AU=WONG JAMES F  
 E7 4 AU=WONG JAMES F H  
 E8 2 AU=WONG JAMES G  
 E9 1 AU=WONG JAMES K  
 E10 12 AU=WONG JAMES M  
 E11 1 AU=WONG JAMES MIN-LEONG  
 E12 14 AU=WONG JAMES R

S36 51 E3,E6-E7  
 S37 410 S35 OR S36  
 S38 339 S37 AND (S1 OR S17)

Ref Items Index-term  
 E1 6 AU=LU AIMING  
 E2 1 AU=LU AINI  
 E3 1 AU=LU AINIAN  
 E4 14 AU=LU AIPIN  
 E5 8 AU=LU AIQIN  
 E6 2 AU=LU AIWU

S30 219 E3, E11

S39 338 S38 AND S2  
S40 338 ID (sorted in duplicate order)  
S41 3201080 DNA OR GENE OR PLASMID  
S42 33 S39 AND S41  
S43 33 ID (sorted in duplicate order)

- 6/6/28 (Item 28 from file: 5) 0006267342 BIOSIS NO.: 198886107263  
PHARMACOLOGY OF POTASSIUM CHANNELS IN THE PLASMALEMA OF THE GREEN ALGA CHARA-CORALLINA 1988
- 6/6/29 (Item 29 from file: 155) 07773735 PMID: 2433153  
Identification of two toxins from scorpion (*Leiurus quinquestriatus*) venom which block distinct classes of calcium-activated potassium channel  
Dec 1 1986
- 6/6/30 (Item 30 from file: 155) 08722670 PMID: 2476127  
Interactions between dendrotoxin, a blocker of voltage-dependent potassium channels, and charybdotoxin, a blocker of calcium-activated potassium channels, at binding sites on neuronal membranes. Aug 30 1989
- 6/6/31 (Item 31 from file: 5) 0006799478 BIOSIS NO.: 198988114593  
INTERACTION BETWEEN DENDROTOXIN A BLOCKER OF VOLTAGE-DEPENDENT POTASSIUM CHANNELS AND CHARYBDOTOXIN A BLOCKER OF CALCIUM-ACTIVATED POTASSIUM CHANNELS AT BINDING SITES ON NEURONAL MEMBRANES 1989
- 6/6/32 (Item 32 from file: 5) 0014988561 BIOSIS NO.: 20040035350  
Kbt1, a three disulfide bridges toxin from *Buthus occitanus tunetanus* venom highly active on both SK and Kv channels 2004
- 6/6/33 (Item 33 from file: 5) 0006395808 BIOSIS NO.: 198936104699  
LEIURUS-QUINQUESTRIATUS VENOM PEPTIDES THAT BLOCK BRAIN VOLTAGE-GATED AND CALCIUM-ACTIVATED POTASSIUM CHANNELS ALSO INHIBIT DENDROTOXIN BINDING TO SYNAPTIC MEMBRANES 1989
- 6/6/34 (Item 34 from file: 155) 10823421 PMID: 7819188  
NMR sequential assignments and solution structure of chlorotoxin, a small scorpion toxin that blocks chloride channels. Jan 10 1995
- 6/6/35 (Item 35 from file: 5) 0009648827 BIOSIS NO.: 199598116680  
NMR Sequential Assignments and Solution Structure of Chlorotoxin, a Small Scorpion Toxin That Blocks Chloride Channels 1995
- 6/6/36 (Item 36 from file: 155) 10895806 PMID: 7533951  
Neuromuscular effects of some potassium channel blocking toxins from the venom of the scorpion *Leiurus quinquestriatus* hebreus. Nov 1994
- 6/6/37 (Item 37 from file: 5) 0009575215 BIOSIS NO.: 199598043048  
Neuromuscular effects of some potassium channel blocking toxins from the venom of the scorpion *Leiurus quinquestriatus* hebreus 1994
- 6/6/38 (Item 38 from file: 155) 09990086 PMID: 1280139  
Noxiustoxin and leiurotoxin III, two homologous peptide toxins with binding properties to synaposomal membrane K<sup>+</sup> channels. Sep 1992
- 6/6/39 (Item 39 from file: 5) 0008709107 BIOSIS NO.: 199395011373  
Noxiustoxin and leiurotoxin III, two homologous peptide toxins with binding properties to synaposomal membrane potassium channels 1992
- 6/6/40 (Item 40 from file: 155) 06794986 PMID: 6197125  
A study on the membrane depolarization of skeletal muscles caused by a scorpion toxin, sea anemone toxin II and crotamine and the interaction between toxins. Jul 1983
- 6/6/41 (Item 41 from file: 5) 0007309715 BIOSIS NO.: 199090094194  
POLARIZED RUBIDIUM-86 EFFLUXES IN PRIMARY CULTURES OF RABBIT KIDNEY PROXIMAL CELLS ROLE OF CALCIUM AND HYPOTONICITY 1990
- 6/6/42 (Item 42 from file: 155) 09066202 PMID: 2165808  
Polarized 86Rb<sup>+</sup> effluxes in primary cultures of rabbit kidney proximal cells: role of calcium and hypotonicity. Jul 9 1990
- 6/6/43 (Item 43 from file: 5) 0005550519 BIOSIS NO.: 198783029410  
PURIFICATION OF CHARYBDOTOXIN A SPECIFIC INHIBITOR OF THE HIGH-CONDUCTANCE CALCIUM-ACTIVATED POTASSIUM CHANNEL 1986
- 6/6/44 (Item 44 from file: 155) 07702384 PMID: 2429958  
Purification of charybdotoxin, a specific inhibitor of the high-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channel Nov 5 1986
- 6/6/45 (Item 45 from file: 155) 08257702 PMID: 2453055  
Purification, sequence, and model structure of charybdotoxin, a potent selective inhibitor of calcium-activated potassium channels. May 1988
- 6/6/46 (Item 46 from file: 5) 0006204047 BIOSIS NO.: 198886043988  
PURIFICATION SEQUENCE AND MODEL STRUCTURE OF CHARYBDOTOXIN A POTENT SELECTIVE INHIBITOR OF CALCIUM-ACTIVATED POTASSIUM CHANNELS 1988
- 6/6/47 (Item 47 from file: 5) 0009196865 BIOSIS NO.: 199497218150  
Solution structure of a core peptide derived from scyllatoxin 1994
- 6/6/48 (Item 48 from file: 155) 13306597 PMID: 10081954  
Solution structure of potassium channel-inhibiting scorpion toxin Lq2. Mar 1 1999
- 6/6/49 (Item 49 from file: 5) 0008984542 BIOSIS NO.: 199497005827  
Synthesis of charybdotoxin and of two N-terminal truncated analogues: Structural and functional characterization 1993
- 6/6/50 (Item 50 from file: 5) 0007639797 BIOSIS NO.: 199191022688  
SYNTHESIS AND STRUCTURAL CHARACTERIZATION OF CHARYBDOTOXIN A POTENT PEPTIDYL INHIBITOR OF THE HIGH CONDUCTANCE CALCIUM ION ACTIVATED POTASSIUM ION CHANNEL 1990
- 6/6/51 (Item 51 from file: 155) 09151531 PMID: 1699936  
Synthesis and structural characterization of charybdotoxin, a potent peptidyl inhibitor of the high conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channel  
Nov 5 1990
- 6/6/52 (Item 52 from file: 5) 0010206293 BIOSIS NO.: 199698674126  
Synthesis and structural characterization of analogues of the potassium channel blocker charybdotoxin 1996
- 6/6/53 (Item 53 from file: 5) 0009025088 BIOSIS NO.: 199497046373  
Toxin pharmacology of the large-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channel in the apical membrane of rabbit proximal convoluted tubule in primary culture 1993
- 6/6/54 (Item 54 from file: 155) 10398153 PMID: 7505914  
Toxin pharmacology of the large-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channel in the apical membrane of rabbit proximal convoluted tubule in primary culture. Oct 1993
- 6/6/55 (Item 55 from file: 155) 09744035 PMID: 1373656  
Toxin pharmacology of the ATP-induced hyperpolarization in Madin-Darby canine kidney cells. Mar 23 1992
- 6/6/56 (Item 56 from file: 5) 0008332816 BIOSIS NO.: 199294034657  
TOXIN PHARMACOLOGY OF ATP-INDUCED HYPERPOLARIZATION IN MADIN-DARBY CANINE KIDNEY CELLS 1992
- 6/6/57 (Item 57 from file: 5) 0006381782 BIOSIS NO.: 198936090673  
TOXINS IN THE CHARACTERIZATION OF POTASSIUM CHANNELS 1989
- 6/6/58 (Item 58 from file: 155) 10066281 PMID: 7678959  
Toxin sensitivity of the calcium-dependent rubidium efflux in Madin-Darby canine kidney cells. Jan 29 1993
- 6/6/59 (Item 59 from file: 5) 0008801098 BIOSIS NO.: 199395103384  
Toxin sensitivity of the calcium-dependent rubidium efflux in Madin-Darby canine kidney cells 1993
- 6/6/60 (Item 60 from file: 155) 08833309 PMID: 2600838  
A voltage-dependent outward current with fast kinetics in single smooth muscle cells isolated from rabbit portal vein. May 1989
- 6/6/61 (Item 61 from file: 5) 0006708256 BIOSIS NO.: 198988023371  
A VOLTAGE-DEPENDENT OUTWARD CURRENT WITH FAST KINETICS IN SINGLE SMOOTH MUSCLE CELLS ISOLATED FROM RABBIT PORTAL VEIN 1989
- 6/6/62 (Item 62 from file: 5) 0007742476 BIOSIS NO.: 199191125367  
THREE-DIMENSIONAL STRUCTURE OF NATURAL CHARYBDOTOXIN IN AQUEOUS SOLUTION BY PROTON NMR CHARYBDOTOXIN POSSESSES A STRUCTURAL MOTIF FOUND IN OTHER SCORPION TOXINS 1991
- 6/7/2 (Item 2 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.  
08751880 PMID: 2477548  
Analysis of the blocking activity of charybdotoxin homologs and iodinated derivatives against Ca<sup>2+</sup>-activated K<sup>+</sup> channels.  
Lucchesi K; Ravindran A; Young H; Moczydlowski E  
Department of Pharmacology, Yale University School of Medicine, New Haven, Connecticut 06510.  
Journal of membrane biology (UNITED STATES) Aug 1989; 109 (3) p269-81; ISSN 0022-2631 Journal Code: 0211301 Contract/Grant No.: AR38796; AR; NIAMS; HL38156; HL, NHLBI Publishing Model Print  
Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM Record type: MEDLINE; Completed  
Two charybdotoxin peptides were purified from venom of the Israeli scorpion, *Leiurus quinquestriatus* hebraeus. Microsequencing of the most abundant toxin, ChTX-Lq1, revealed identity with the 37-residue peptide previously sequenced by Gimenez-Gallego et al. [Gimenez-Gallego, G., et al., Proc. Natl. Acad. Sci. USA 85:3329-3333 (1988)]. Sequence data on the minor peptide, ChTX-Lq2, showed substantial homology to ChTX-Lq1 with differences observed at eight positions. These two charybdotoxin sequences, along with that of noxiustoxin, define a distinct family of scorpion peptide toxins with activity against K<sup>+</sup> channels. Both charybdotoxin homologs inhibited Ca<sup>2+</sup>-dependent K<sup>+</sup> efflux from human erythrocytes with similar potency, K<sub>0.5</sub> approximately 40 nM. In planar bilayer assays of single K(Ca) channels from rat muscle, ChTX-Lq1 and ChTX-Lq2 blocked with intrinsic K<sub>d</sub>s of 1.3 and 43 nM, respectively, in the presence of 50 mM external KCl. A new application of dwell-time histogram analysis of single-channel blocking events was used to characterize the kinetic homogeneity of toxin samples and the blocking kinetics of ChTX derivatives. The lower blocking affinity of ChTX-Lq2

was the combined result of a faster dissociation rate and a slower association rate as compared to ChTX-Lq1. The blocking activity of two mono-iodinated derivatives of ChTX-Lq1 was also analyzed. Blocked dwell-time histograms of the iodinated peptides were characterized by predominantly brief (0.2-2 sec) blocking events in comparison to the native toxin (20 sec). Histogram analysis revealed that mono-iodination of ChTX-Lq1 impairs blocking activity by adverse effects on both dissociation and association rate constants. Frequency density histograms of single channel blocking events provide a sensitive assay of toxin purity suitable for quantitating structure-activity relationships of charybdotoxin derivatives.

Record Date Created: 19891101 Record Date Completed: 19891101

6/7/6 (Item 6 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.  
08675595 PMID: 2473920

Charybdotoxin blocks both Ca-activated K channels and Ca-independent voltage-gated K channels in rat brain synaptosomes.

Schneider M J; Rogowski R S; Krueger B K; Blaustein M P  
Department of Physiology, University of Maryland School of Medicine, Baltimore 21201.

FEBS letters (NETHERLANDS) Jul 3 1989; 250 (2) p433-6; ISSN 0014-5793 Journal Code: 0155157

Contract/Grant No.: NS-16106; NS; NINDS; NS-16285; NS; NINDS; NS-20106; NS; NINDS Publishing Model Print Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM Record type: MEDLINE; Completed

Charybdotoxin (ChTX), a 4.3 kDa polypeptide toxin from the venom of the scorpion *Leiurus quinquestriatus*, blocks both a Ca-activated K channel (IC50 approximately 15 nM) and a Ca-independent voltage-gated K channel (IC50 approximately 40 nM) in rat brain synaptosomes. These results indicate that in this preparation ChTX is not specific for the Ca-activated K channel and suggest that there may be structural homology among the toxin-binding sites on various types of K channels.

Record Date Created: 19890907 Record Date Completed: 19890907

6/7/9 (Item 9 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.  
11058434 PMID: 7543240

Charybdotoxin and its effects on potassium channels.

Garcia M L; Kraus H G; Munujos P; Slaughter R S; Kaczorowski G J

Department of Membrane Biochemistry and Biophysics, Merck Research Laboratories, Rahway, New Jersey 07065, USA. American journal of physiology (UNITED STATES) Jul 1995; 269 (1 Pt 1) pC1-10; ISSN 0002-9513 Journal Code: 0370511 Publishing Model Print Document type: Journal Article; Review; Tutorial Languages: ENGLISH

Main Citation Owner: NLM Record type: MEDLINE; Completed

Over the last few years, a considerable amount of information has been obtained regarding K+ channels. Different areas of research have contributed to knowledge in this field. Charybdotoxin (ChTX), a 37-amino acid peptide isolated from venom of the scorpion *Leiurus quinquestriatus* var. *hebraeus*, represents a remarkable tool for studying K+ channels. With its use, it has been possible to purify the high-conductance Ca(2+)-activated K+ (maxi-K) channel to homogeneity and determine the subunit composition of this channel. This has led to the discovery of an auxiliary beta-subunit that, when coexpressed with the pore-forming subunit, mSlo, alters the biophysical and pharmacological properties of this latter subunit. With the feasibility of producing large amounts of ChTX by recombinant techniques and the knowledge of the three-dimensional structure of the peptide, it has been possible to carry out site-directed mutagenesis studies and obtain a picture of the interaction surface of the toxin with two channels, maxi-K and Shaker, and to derive a picture of the complementary, surface of the receptor in these two channels. Finally, ChTX, and the more selective K+ channel toxins that were subsequently discovered, have provided us with unique tools not only to determine the functional role that K+ channels play in target tissues but also to develop the molecular pharmacology of these channels. (76 Refs.)

Record Date Created: 19950901 Record Date Completed: 19950901

6/7/11 (Item 11 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.  
08859199 PMID: 2482078

Charybdotoxin is a new member of the K+ channel toxin family that includes dendrotoxin I and mast cell degranulating peptide.

Schweitz H; Bidaud J N; Maes P; Lazdunski M

Centre de Biochimie, Centre National de la Recherche Scientifique, Université de Nice, France.

Biochemistry (UNITED STATES) Dec 12 1989; 28 (25) p9708-14; ISSN 0006-2960 Journal Code: 0370623

Publishing Model Print Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM

Record type: MEDLINE; Completed

A polypeptide was identified in the venom of the scorpion *Leiurus quinquestriatus hebraeus* by its potency to inhibit the high-affinity binding of the radiolabeled snake venom toxin dendrotoxin I (125I-DTX1) to its receptor site. It has been purified, and its properties investigated by different techniques were found to be similar to those of MCD and DTX1, two polypeptide toxins active on a voltage-dependent K+ channel. However, its amino acid sequence was determined, and it was shown that this toxin is in fact charybdotoxin (ChTX), a toxin classically used as a specific tool to block one class of Ca2+-activated K+ channels. ChTX, DTX1, and MCD are potent convulsants and are highly toxic when injected

intracerebroventricularly in mice. Their toxicities correlate well with their affinities for their receptors in rat brain. These three structurally different toxins release [3H]GABA from preloaded synaptosomes, the efficiency order being DTX1 greater than ChTX greater than MCD. Both binding and cross-linking experiments of ChTX to rat brain membranes and to the purified MCD/DTX1 binding protein have shown that the alpha-subunit (Mr = 76K-78K) of the MCD/DTX1-sensitive K+ channel protein also contains the ChTX binding sites. Binding sites for DTX1, MCD, and ChTX are in negative allosteric interaction. Our results show that charybdotoxin belongs to the family of toxins which already includes the dendrotoxins and MCD, which are blockers of voltage-sensitive K+ channels. ChTX is clearly not selective for Ca2+-activated K+ channel.

Record Date Created: 19900305 Record Date Completed: 19900305

6/7/18 (Item 18 from file: 5) DIALOG(R)File 5:Biosis Previews(R) (c) 2005 BIOSIS. All rts. reserv.  
0011349339 BIOSIS NO.: 199800143586

Consequence of the removal of evolutionary conserved disulfide bridges on the structure and function of charybdotoxin and evidence that particular cysteine spacing govern specific disulfide bond formation

AUTHOR: Drakopoulou Eugenia; Vizzanova Jean; Neyton Jacques; Anriot Vincent ; Bouet Franciose; Virelizier Henri; Menez Andre; Vita Claudio (Reprint)

AUTHOR ADDRESS: CEA, Dep. Ingeniere Etudes Proteines, Serv. Phys. Exp. Analyse, Saclay, 91190 Gif-sur-Yvette, France\*\*France

JOURNAL: Biochemistry 37 (5); p1292-1301 Feb. 3, 1998 1998 MEDIUM: print ISSN: 0006-2960 DOCUMENT TYPE: Article RECORD TYPE: AbstractLANGUAGE: English

ABSTRACT: Scorpion toxins are miniglobular proteins containing a common structural motif formed by an alpha-helix on one face, an antiparallel beta-sheet on the opposite face, and three disulfide bonds making up most of its internal volume. We have investigated the role of these evolutionary conserved bonds by replacing each couple of bridged cysteine residues of the scorpion charybdotoxin by a pair of nonbridging L-alpha-aminobutyric acid (Abu) residues. Three analogues were obtained by solid-phase synthesis, Chab I, Chab II, and Chab III, containing the Abu residues in positions 7 and 28, 13 and 33, 17 and 35, respectively. Circular dichroism analysis showed that the purified Chab II acquired a conformation similar to that of charybdotoxin, while the Chab I and Chab III possess decreased native-like characteristics. All analogues block single high-conductance Ca2+-activated K+ channels from rat skeletal muscle inserted into planar lipid bilayers, but with different potencies. Chab II is the most active analogue (KD = 8.0 X 10-8 M), with a 9-fold lower affinity as compared to native charybdotoxin. Chab I and Chab III have, respectively, 180- and 580-fold lower affinity. Therefore, the removal of evolutionary conserved disulfide bridges does not prevent the toxin to adopt a functional and presumably native-like structure. However, removal of one disulfide bond affects the yields of formation of correct pairing between the remaining cysteine residues, and only Chab I preserves the ability to form the native disulfide pairings with high efficiency. This is the only analogue to preserve particular spacings of three and one residue between the cysteines, which have been described to thermodynamically disfavor disulfide bond formation between the cysteines (Zhang, R., and Snyder, G. H. (1989) J. Biol. Chem. 264, 18472-18479). Therefore, we conclude that the position of the cysteine residues in the sequence of charybdotoxin, by disfavoring specific pairings and favoring others, may govern selective formation of specific disulfide bonds, thus, explaining the efficient folding properties of Chab I and of native charybdotoxin. The structural properties of the Chab analogues and the discovered role of the cysteine spacings have interesting implications in protein design and engineering.

6/7/19 (Item 19 from file: 5) DIALOG(R)File 5:Biosis Previews(R) (c) 2005 BIOSIS. All rts. reserv.  
0007735556 BIOSIS NO.: 19919118447

DESIGN SYNTHESIS AND FUNCTIONAL EXPRESSION OF A GENE FOR CHARYBDOTOXIN A PEPTIDE BLOCKER OF POTASSIUM ION CHANNELS

AUTHOR: PARK C-S (Reprint); HAUSDORFF S F; MILLER C

AUTHOR ADDRESS: HOWARD HUGHES MED INST GRADUATE DEP BIOCHEM, BRANDEIS UNIV, WALTHAM, MASS 02254, USA\*\*USA

JOURNAL: Proceedings of the National Academy of Sciences of the United States of America 88 (6); p2046-2050 1991 ISSN: 0027-8424 DOCUMENT TYPE: Article RECORD TYPE: Abstract LANGUAGE: ENGLISH

ABSTRACT: A gene encoding charybdotoxin (CTX), a K+ channel blocker from scorpion venom, was designed, synthesized, and expressed as a cleavable fusion protein in *Escherichia coli*. A sequence-specific protease, factor Xa, was used to cleave the fusion protein and thus release the toxin peptide. The recombinant toxin was purified, oxidized to form disulfide bonds, and treated to form N-terminal pyroglutamate. Recombinant CTX is identical to the native venom CTX with respect to high-performance liquid chromatography mobility, amino acid composition, and N-terminal modification. With single Ca2+-activated K+ channels as an assay system, recombinant CTX shows blocking and dissociation kinetics identical to the native venom toxin. The synthetic gene and high-level expression of functionally active CTX make it possible to study the fundamental mechanism of the toxin-ion channel interaction.

6/7/22 (Item 22 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.

08812992 PMID: 2531622

Effects of potassium channel toxins from *Leiurus quinquestriatus hebraeus* venom on responses to cromakalim in rabbit blood vessels.

Strong P N; Weir S W; Beech D J; Hiestand P; Kocher H P

Jerry Lewis Muscle Research Centre, Department of Paediatrics and Neonatal Medicine, Royal Postgraduate Medical School, London.

British journal of pharmacology (ENGLAND) Nov 1989, 98 (3) p817-26, ISSN 0007-1188 Journal Code: 7502536  
 Publishing Model Print Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM  
 Record type: MEDLINE; Completed

1. The effects of fractionated Leiurus quinquestriatus hebraeus venom on cromakalim-induced 86Rb<sup>+</sup> efflux in rabbit aortic smooth muscle were examined. 2. Crude venom (0.1-30 micrograms ml<sup>-1</sup>) produced a concentration-dependent decrease of 1 microM cromakalim-induced 86Rb<sup>+</sup> response. The maximum blocking activity attainable was approximately 60%. 3. Fractionation of crude venom by gel permeation chromatography and subsequent chromatography on a cation ion-exchange column, produced two fractions (X and XI), active in the 86Rb<sup>+</sup> blocking assay. 4. Fraction XII contained charybdotoxin (approximately 85% pure). After a final high performance liquid chromatography (h.p.l.c.) purification step, the purified toxin failed to inhibit the cromakalim-stimulated 86Rb<sup>+</sup> efflux although it was a potent inhibitor of A23187-induced K<sup>+</sup> flux in human erythrocytes and the large conductance calcium-activated potassium channel in rabbit portal vein smooth muscle. 5. Subsequent purification of fraction X by h.p.l.c. yielded a minor peak which contained 86Rb<sup>+</sup> blocking activity. This subfraction was also capable of inhibiting apamin-sensitive, angiotensin II-stimulated K<sup>+</sup> flux in guinea-pig hepatocytes. 6. It is concluded that the potassium channel opened by cromakalim in rabbit aortic smooth muscle is not blocked by charybdotoxin but by another distinct toxin in the venom of Leiurus quinquestriatus hebraeus.

Record Date Created: 19900125 Record Date Completed: 19900125

6/7/29 (Item 29 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.  
 07773735 PMID: 2433153  
 Identification of two toxins from scorpion (Leiurus quinquestriatus) venom which block distinct classes of calcium-activated potassium channel.  
 Castle N A; Strong P N  
 FEBS letters (NETHERLANDS) Dec 1 1986, 209 (1) p117-21, ISSN 0014-5793 Journal Code: 0155157  
 Publishing Model Print Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM  
 Record type: MEDLINE; Completed

Two polypeptide toxins from scorpion (Leiurus quinquestriatus) venom which block distinct classes of calcium-activated potassium channels have been identified and partially purified. One toxin, at 50-100 ng/ml, blocks apamin-sensitive potassium fluxes in hepatocytes and inhibits [125I]moroindopamine binding. The other, more basic, toxin blocks apamin-insensitive potassium fluxes in erythrocytes at 200 ng/ml and, to our knowledge, is the first toxin shown to block the erythrocyte calcium-activated potassium channel with high affinity. The possible co-identity of this latter toxin with charybdotoxin is discussed.

Record Date Created: 19870220 Record Date Completed: 19870220

6/7/33 (Item 33 from file: 5) DIALOG(R)File 5:Biosis Previews(R) (c) 2005 BIOSIS. All rts. reserv.  
 0006395808 BIOSIS NO.: 198936104699  
 LEIURUS- QUINQUESTRIATUS VENOM PEPTIDES THAT BLOCK BRAIN VOLTAGE-GATED AND CALCIUM-ACTIVATED POTASSIUM CHANNELS ALSO INHIBIT DENDROTOXIN BINDING TO SYNAPTIC MEMBRANES  
 AUTHOR: SORENSSEN R G (Reprint); SCHNEIDER M J; BLAUSTEIN M P  
 AUTHOR ADDRESS: MED DEP, DIV ENVIRON MED AND TOXICOL, JEFFERSON MED COLL, PHILADELPHIA, PA 19107, USA\*\*USA  
 JOURNAL: Biophysical Journal 55 (2 PART 2): p550A 1989 CONFERENCE/MEETING: THIRTY-THIRD ANNUAL MEETING OF THE BIOPHYSICAL SOCIETY, CINCINNATI, OHIO, USA, FEBRUARY 12-16, 1989 BIOPHYS J ISSN: 0006-3495  
 DOCUMENT TYPE: Meeting RECORD TYPE: Citation LANGUAGE: ENGLISH

6/7/36 (Item 36 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.  
 10895806 PMID: 7533951  
 Neuromuscular effects of some potassium channel blocking toxins from the venom of the scorpion Leiurus quinquestriatus hebraeus.  
 Marshall D L; Vatanpour H; Harvey A L; Boyot P; Pinkasfeld S; Doljansky Y; Bouel F; Menez A  
 Department of Physiology and Pharmacology, University of Strathclyde, Glasgow, U.K.  
 Toxicon - official journal of the International Society on Toxicology (ENGLAND) Nov 1994, 32 (11) p1433-43, ISSN 0041-0101 Journal Code: 1307333 Publishing Model Print Document type: Journal Article Languages: ENGLISH  
 Main Citation Owner: NLM Record type: MEDLINE; Completed

The scorpion venom Leiurus quinquestriatus hebraeus was fractionated by chromatography in order to isolate toxins that affected binding of radiolabelled dendrotoxin to K<sup>+</sup> channel proteins on synaptosomal membranes and that facilitated acetylcholine release in chick biventer cervicis nerve-muscle preparations. In addition to the previously characterized charybdotoxin, three toxins were isolated: 14-2, 15-1 and 18-2. Toxin 14-2 has a blocked N-terminus and because of low quantities, it has not been sequenced; 15-1 is a newly sequenced toxin of 36 residues with some overall homology to charybdotoxin and noxiustoxin; 18-2 is identical to charybdotoxin -2. The apparent K<sub>i</sub> against dendrotoxin binding were: charybdotoxin, 3.8 nM; 14-2, 150 nM; 15-1, 50 nM; and 18-2, 0.25 nM. Toxin 14-2 (75 nM-1.5 microm) had a presynaptic

facilitatory effect on neuromuscular preparations. Toxin 15-1 augmented responses to direct muscle stimulation, probably because it blocked Ca<sup>2+</sup>-activated K<sup>+</sup> currents in muscle fibres. Toxin 18-2 (charybdotoxin -2) had a potent presynaptic facilitatory action, with less effect on direct muscle stimulation. This contrasts with the relatively weak neuromuscular effects of the highly homologous charybdotoxin. On a Ca<sup>2+</sup>-activated K<sup>+</sup> current in mouse motor nerve endings, charybdotoxin and toxin 18-2 produced maximal block at around 100 nM, whereas 15-1 was inactive at 300 nM. Charybdotoxin can increase quantal content, but this is more likely to result from block of voltage-dependent K<sup>+</sup> channels than Ca<sup>2+</sup>-activated channels: the increase in transmitter release occurred in conditions in which little IKCa would be present; higher concentration of charybdotoxin and longer exposure times were required to increase transmitter release than those needed to block IKCa, and the facilitatory effects of charybdotoxin and toxin 18-2 correlated more with their effects on dendrotoxin binding than on block of IKCa.

Record Date Created: 19950410 Record Date Completed: 19950410

6/7/45 (Item 45 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.  
 08257702 PMID: 2453055  
 Purification, sequence, and model structure of charybdotoxin, a potent selective inhibitor of calcium-activated potassium channels.  
 Gimenez-Gallego G; Navia M A; Reuben J P; Katz G M; Kaczorowski G J; Garcia M L  
 Department of Growth Factor Research, Merck Sharp & Dohme Research Laboratories, Rahway, NJ 07065.  
 Proceedings of the National Academy of Sciences of the United States of America (UNITED STATES) May 1988, 85 (10) p3329-33, ISSN 0027-8424 Journal Code: 7505876 Publishing Model Print Document type: Journal Article  
 Languages: ENGLISH Main Citation Owner: NLM Record type: MEDLINE; Completed

Charybdotoxin (ChTX), a protein present in the venom of the scorpion Leiurus quinquestriatus var hebraeus, has been purified to homogeneity by a combination of ion-exchange and reversed-phase chromatography. Polyacrylamide gel electrophoresis, amino acid analysis, and complete amino acid sequence determination of the pure protein reveal that it consists of a single polypeptide chain of 4.3 kDa. Purified ChTX is a potent and selective inhibitor of the approximately 220-pS Ca<sup>2+</sup>-activated K<sup>+</sup> channel present in GH3 anterior pituitary cells and primary bovine aortic smooth muscle cells. The toxin reversibly blocks channel activity by interacting at the external pore of the channel protein with an apparent K<sub>d</sub> of 2.1 nM. The primary structure of ChTX is similar to a number of neurotoxins of diverse origin, which suggests that ChTX is a member of a superfamily of proteins that modify ion-channel activities. On the basis of this similarity, the three-dimensional structure of ChTX has been modeled from the known crystal structure of alpha-bungarotoxin. These studies indicate that ChTX is useful as a probe of Ca<sup>2+</sup>-activated K<sup>+</sup>-channel function and suggest that the proposed tertiary structure of ChTX may provide insight into the mechanism of channel block.

Record Date Created: 19880622 Record Date Completed: 19880622

6/7/52 (Item 52 from file: 5) DIALOG(R)File 5:Biosis Previews(R) (c) 2005 BIOSIS. All rts. reserv.  
 0010206293 BIOSIS NO.: 19969874126  
 Synthesis and structural characterisation of analogues of the potassium channel blocker charybdotoxin  
 AUTHOR: Dyke Timothy R; Duggan Brendan M; Pennington Michael W; Byrnes Michael E; Kem William R; Norton Raymond S (Reprint)  
 AUTHOR ADDRESS: NMR Lab., Biomolecular Research Inst., 381 Royal Parade, Parkville, VIC 3052, Australia\*\*Australia  
 JOURNAL: Biochimica et Biophysica Acta 1292 (1): p31-38 1996 1996 ISSN: 0006-3002 DOCUMENT TYPE: Article  
 RECORD TYPE: Abstract LANGUAGE: English

ABSTRACT: Charybdotoxin is a 37-residue polypeptide toxin from scorpion venom, which acts by blocking voltage-gated and Ca<sup>2+</sup>-activated K<sup>+</sup> channels. We have synthesized charybdotoxin and three mono-substituted analogues using an Fmoc-tBu protocol. The Phe-2 fwdaw Tyr analogue was chosen to introduce a site for Tyr iodination which was distinct from the K<sup>+</sup> channel binding surface, while the Glu-12 fwdaw Gln and Arg-19 fwdaw His analogues were studied to probe the roles of charged residues at these positions in the structure and activity of the toxin. The synthetic native molecule was equipotent with natural toxin in inhibiting the human erythrocyte Ca<sup>2+</sup>-dependent K<sup>+</sup> channel. The affinities of all three analogues for the erythrocyte K<sup>+</sup> channel were slightly reduced, with the Arg-19 fwdaw His analogue showing the greatest increase in IC<sub>50</sub> (2.3-fold). Two-dimensional <sup>1</sup>H-NMR studies of these analogues showed that all three had structures very similar to those of the native molecule. The most significant perturbation was associated with the Glu-12 to Gln substitution, which appeared to destabilise the N-terminal half of the alpha-helix, possibly due to the weakening of an N-terminal helix capping interaction which is apparent from our NMR data. His-21 has a pK<sub>a</sub> more than one unit below the value for a non-interacting histidine. Possible reasons for this are that the imidazolium side chain is partly buried and is located near positively charged moieties. Thus, His-21 would be neutral at physiological pH, where charybdotoxin binds to the potassium channel.

6/7/62 (Item 62 from file: 5) DIALOG(R)File 5:Biosis Previews(R) (c) 2005 BIOSIS. All rts. reserv.  
 0007742476 BIOSIS NO.: 199191125367  
 THREE-DIMENSIONAL STRUCTURE OF NATURAL CHARYBDOTOXIN IN AQUEOUS SOLUTION BY PROTON NMR  
 CHARYBDOTOXIN POSSESSES A STRUCTURAL MOTIF FOUND IN OTHER SCORPION TOXINS  
 AUTHOR: BONTEMPS F (Reprint); ROUMESTAND C; BOYOT P; GILQUIN B; DOLJANSKY Y; MENEZ A; TOMA F

**AUTHOR ADDRESS:** SERV DE BIOCHIM DES PROTEINES, LAB D'INGENIERIE DES PROTEINES BAT 152, CEN-SACLAY, F-91191 GIF-SUR-YVETTE, FR\*\*FRANCE  
**JOURNAL:** European Journal of Biochemistry 196 (1): p19-28 1991 ISSN: 0014-2956 DOCUMENT TYPE: Article  
**RECORD TYPE:** Abstract LANGUAGE: ENGLISH  
**ABSTRACT:** A 600-MHz proton NMR study of the scorpion toxin k $\kappa$ -conotoxin PVIIA, a toxin acting on K $^{+}$  channels, is reported. The unambiguous sequential assignment of all the protons of the toxin was achieved. The analysis of NOEs and of backbone coupling constants showed the existence of an  $\alpha$ -helix (residues 10-19) and of an antiparallel  $\beta$ -sheet in the 26-35 part. Three-dimensional structures were generated by distance geometry, using a set of 114 inter-residual calibrated constraints (63 sequential, 47 medium and long range, 4 hydrogen bonds) and 29  $\phi$ ,  $\psi$  angles. These structures show that charybotoxin is composed of a  $\beta$ -sheet linked to an  $\alpha$ -helix by two disulphide bridges and to an extended fragment by the third disulphide bridge. Comparison with the other known structures of long and short scorpion toxins shows that this structural motif is common to all these proteins.

- 9/6/1 (Item 1 from file: 5) 0007115178 BIOSIS NO.: 1990890330069  
 ANALYSIS OF THE BLOCKING ACTIVITY OF CHARYBOTOXIN HOMOLOGS AND IODINATED DERIVATIVES AGAINST CALCIUM-ACTIVATED POTASSIUM CHANNELS 1989
- 9/6/2 (Item 2 from file: 155) 08751880 PMID: 2477548  
 Analysis of the blocking activity of charybotoxin homologs and iodinated derivatives against Ca $^{2+}$ -activated K $^{+}$  channels. Aug 1989
- 9/6/3 (Item 3 from file: 155) 08575595 PMID: 2473920  
 Charybotoxin blocks both Ca-activated K channels and Ca-independent voltage-gated K channels in rat brain synaptosomes. Jul 3 1989
- 9/6/4 (Item 4 from file: 155) 0859199 PMID: 2482078  
 Charybotoxin is a new member of the K $^{+}$ -channel toxin family that includes dendrotoxin I and mast cell degranulating peptide. Dec 12 1989
- 9/6/5 (Item 5 from file: 5) 0007133033 BIOSIS NO.: 199089050924  
 CHARYBOTOXIN IS A NEW MEMBER OF THE POTASSIUM CHANNEL TOXIN FAMILY THAT INCLUDES DENDROTOXIN I AND MAST CELL DEGRANULATING PEPTIDE 1989
- 9/6/6 (Item 6 from file: 5) 0006941512 BIOSIS NO.: 199038119403  
 CHARYBOTOXIN IS A NEW MEMBER OF THE TOXIN FAMILY THAT INCLUDES DENDROTOXIN I AND MCD AND BLOCKS DENDROTOXIN-SENSITIVE VOLTAGE-ACTIVATED POTASSIUM CHANNELS 1990
- 9/6/7 (Item 7 from file: 5) 0011349339 BIOSIS NO.: 199800143586  
 Consequence of the removal of evolutionary conserved disulfide bridges on the structure and function of charybotoxin and evidence that particular cysteine spacing govern specific disulfide bond formation 1998
- 9/6/8 (Item 8 from file: 155) 08571405 PMID: 2468961  
 Effect of some potassium channel blockers on contractile responses of the rabbit aorta. Feb 1989
- 9/6/9 (Item 9 from file: 5) 0006395808 BIOSIS NO.: 198936104699  
 LEIURUS-QUINQUESTRIATUS VENOM PEPTIDES THAT BLOCK BRAIN VOLTAGE-GATED AND CALCIUM-ACTIVATED POTASSIUM CHANNELS ALSO INHIBIT DENDROTOXIN BINDING TO SYNAPTIC MEMBRANES 1989
- 9/6/10 (Item 10 from file: 155) 10823421 PMID: 7819188  
 NMR sequential assignments and solution structure of chibrotxin, a small scorpion toxin that blocks chloride channels. Jan 10 1995
- 9/6/11 (Item 11 from file: 5) 0009648827 BIOSIS NO.: 199598116660  
 NMR Sequential Assignments and Solution Structure of Chibrotxin, a Small Scorpion Toxin That Blocks Chloride Channels 1995
- 9/6/12 (Item 12 from file: 155) 10895806 PMID: 7533951  
 Neuromuscular effects of some potassium channel blocking toxins from the venom of the scorpion Leiurus quinquestriatus hebreus. Nov 1994
- 9/6/13 (Item 13 from file: 5) 0009575215 BIOSIS NO.: 199598043048  
 Neuromuscular effects of some potassium channel blocking toxins from the venom of the scorpion Leiurus quinquestriatus hebreus 1994
- 9/6/14 (Item 14 from file: 155) 09990086 PMID: 1280139  
 Noxiustoxin and leiurotoxin III, two homologous peptide toxins with binding properties to synaptosomal membrane K $^{+}$  channels. Sep 1992
- 9/6/15 (Item 15 from file: 5) 0008709107 BIOSIS NO.: 199395011373  
 Noxiustoxin and leiurotoxin III, two homologous peptide toxins with binding properties to synaptosomal membrane potassium channels 1992
- 9/6/16 (Item 16 from file: 5) 0008984542 BIOSIS NO.: 199497005827  
 Synthesis of charybotoxin and of two N-terminal truncated analogues: Structural and functional characterization 1993
- 9/6/17 (Item 17 from file: 5) 0010206293 BIOSIS NO.: 199698674126  
 Synthesis and structural characterisation of analogues of the potassium channel blocker charybotoxin 1996

- 9/6/18 (Item 18 from file: 5) 0009025088 BIOSIS NO.: 199497046373  
 Toxin pharmacology of the large-conductance Ca $^{2+}$ -activated K $^{+}$  channel in the apical membrane of rabbit proximal convoluted tubule in primary culture 1993
- 12/6/1 (Item 1 from file: 155) 11322494 PMID: 8645186  
 A novel structural class of K $^{+}$ -channel blocking toxin from the scorpion Pandinus imperator. May 1 1996
- 13/6/1 (Item 1 from file: 155) 11322494 PMID: 8645186  
 A novel structural class of K $^{+}$ -channel blocking toxin from the scorpion Pandinus imperator. May 1 1996
- 13/6/2 (Item 1 from file: 5) 0010380507 BIOSIS NO.: 199699014567  
 A novel structural class of K $^{+}$ -channel blocking toxin from the scorpion Pandinus imperator 1996
- 16/6/1 (Item 1 from file: 155) 12140757 PMID: 9438859  
 Solution structure and proposed binding mechanism of a novel potassium channel toxin kappa-conotoxin PVIIA. Dec 15 1997
- 16/6/2 (Item 1 from file: 5) 0011233829 BIOSIS NO.: 199800088076  
 Solution structure and proposed binding mechanism of a novel potassium channel toxin kappa-conotoxin PVIIA 1997
- 19/6/1 (Item 1 from file: 155) 10204659 PMID: 7685635  
 An activator of calcium-dependent potassium channels isolated from a medicinal herb. Jun 22 1993
- 19/6/2 (Item 2 from file: 155) 15271073 PMID: 15051409  
 Antigenic polymorphism of the "short" scorpion toxins able to block K $^{+}$  channels. Mar 15 2004
- 19/6/3 (Item 3 from file: 5) 0005645783 BIOSIS NO.: 198783124674  
 BLOCKING AGENTS OF CALCIUM-ACTIVATED POTASSIUM CHANNELS IN CULTURED MEDULLARY THICK ASCENDING LIMB CELLS 1987
- 19/6/4 (Item 4 from file: 155) 07821071 PMID: 2435161  
 Blocking agents of Ca $^{2+}$ -activated K $^{+}$  channels in cultured medullary thick ascending limb cells. Feb 1987
- 19/6/5 (Item 5 from file: 155) 17947714 PMID: 15695820  
 BmP09, a "long chain" scorpion peptide blocker of BK channels. Apr 15 2005
- 19/6/6 (Item 6 from file: 5) 0008848545 BIOSIS NO.: 199396012961  
 Calcium-activated potassium transport in erythrocytes: Comparison of binding and transport inhibition by scorpion toxins 1993
- 19/6/7 (Item 7 from file: 155) 10517036 PMID: 8297371  
 Chemical synthesis and structure-function studies of margatoxin, a potent inhibitor of voltage-dependent potassium channel in human T lymphocytes. Jan 28 1994
- 19/6/8 (Item 8 from file: 155) 08816237 PMID: 2480347  
 Characterization of high affinity binding sites for charybotoxin in sarcolemmal membranes from bovine aortic smooth muscle. Evidence for a direct association with the high conductance calcium-activated potassium channel. Dec 15 1989
- 19/6/9 (Item 9 from file: 155) 11411894 PMID: 8706835  
 Characterization of a new peptide from Tityus serrulatus scorpion venom which is a ligand of the apamin-binding site. Jul 15 1996
- 19/6/10 (Item 10 from file: 155) 16507366 PMID: 15189853  
 Computational simulations of interactions of scorpion toxins with the voltage-gated potassium ion channel. Jun 2004
- 19/6/11 (Item 11 from file: 5) 0014960511 BIOSIS NO.: 200400331297  
 Computational simulations of interactions of scorpion toxins with the voltage-gated potassium ion channel 2004
- 19/6/12 (Item 12 from file: 155) 12172929 PMID: 9477955  
 Consequence of the removal of evolutionary conserved disulfide bridges on the structure and function of charybotoxin and evidence that particular cysteine spacings govern specific disulfide bond formation. Feb 3 1998
- 19/6/13 (Item 13 from file: 155) 11091649 PMID: 7545007  
 Cross-linking of charybotoxin to high-conductance calcium-activated potassium channels: identification of the covalently modified toxin residue. Aug 29 1995
- 19/6/14 (Item 14 from file: 155) 11196506 PMID: 8531201  
 Ca(2+)-activated K $^{+}$  channels of human and rabbit erythrocytes display distinctive patterns of inhibition by venom peptide toxins. Sep 1995
- 19/6/15 (Item 15 from file: 155) 10145953 PMID: 7682555  
 Ca(2+)-activated K $^{+}$  transport in erythrocytes. Comparison of binding and transport inhibition by scorpion toxins. Apr 25 1993
- 19/6/16 (Item 16 from file: 155) 09358728 PMID: 1706481



Determination of the subunit stoichiometry of a voltage-activated potassium channel. Mar 21 1991

19/6/17 (Item 17 from file: 155) 11224805 PMID: 8527429  
Determination of the three-dimensional solution structure of noxiustoxin: analysis of structural differences with related short-chain scorpion toxins. Dec 26 1995

19/6/18 (Item 18 from file: 155) 09905670 PMID: 1381959  
Determination of the three-dimensional structure of iberitoxin in solution by 1H nuclear magnetic resonance spectroscopy. Sep 8 1992

19/6/19 (Item 19 from file: 155) 12196378 PMID: 9504384  
Effect of Thyus serratus scorpion venom on the rabbit isolated corpus cavernosus and the involvement of NANC nitergic nerve fibres. Feb 1998

19/6/20 (Item 20 from file: 155) 13573819 PMID: 11155206  
Effects of toxins P2 and P3 on human T lymphocyte Kv1.3 channels: the role of G17 and Lys24. Jan 1 2001

19/6/21 (Item 21 from file: 155) 08469848 PMID: 2464066  
An emerging pharmacology of peptide toxins targeted against potassium channels. Oct 1988

19/6/22 (Item 22 from file: 155) 14389721 PMID: 12220678  
Glycine 30 in iberitoxin is a critical determinant of its specificity for maxi-K versus K(V) channels. Sep 11 2002

19/6/23 (Item 23 from file: 155) 13705739 PMID: 11352729  
Interaction of a toxin from the scorpion Thyus serratus with a cloned K<sup>+</sup>-channel from squid (sqKv1A). May 22 2001

19/6/24 (Item 24 from file: 155) 17279256 PMID: 15165720  
Kvot1, a three disulfide bridges toxin from Buthus occitanus tunetanus venom highly active on both SK and Kv channels. Apr 2004

19/6/25 (Item 25 from file: 5) 0008234656 BIOSIS NO.: 199293077547  
KALOTOXIN A NOVEL PEPTIDYL INHIBITOR OF NEURONAL BK-TYPE CALCIUM-ACTIVATED POTASSIUM CHANNELS CHARACTERIZED FROM ANDROCTONUS-MAURETANICUS-MAURETANICUS VENOM 1992

19/6/26 (Item 26 from file: 155) 08629861 PMID: 1730708  
Kalotoxin, a novel peptidyl inhibitor of neuronal BK-type Ca(2+)-activated K<sup>+</sup> channels characterized from Androctonus mauretanicus mauretanicus venom. Jan 25 1992

19/6/27 (Item 27 from file: 155) 10742600 PMID: 7524673  
Kalotoxin (1-37) shows structural differences with related potassium channel blockers. Nov 29 1994

19/6/28 (Item 28 from file: 5) 0009610999 BIOSIS NO.: 199598078832  
Kalotoxin (1-37) shows structural differences with related potassium channel blockers 1994

19/6/29 (Item 29 from file: 155) 09853502 PMID: 1379069  
Mechanism of iberitoxin block of the large-conductance calcium-activated potassium channel from bovine aortic smooth muscle. Jul 28 1992

19/6/30 (Item 30 from file: 155) 12969238 PMID: 10920011  
Mechanisms of neurotoxin action on Shaker potassium channels. Aug 2000

19/6/31 (Item 31 from file: 5) 0008413156 BIOSIS NO.: 199294114997  
MODE OF ACTION OF IBERITOXIN A POTENT BLOCKER OF THE LARGE CONDUCTANCE CALCIUM ACTIVATED POTASSIUM CHANNEL 1992

19/6/32 (Item 32 from file: 155) 09957833 PMID: 1384740  
Mode of action of iberitoxin, a potent blocker of the large conductance Ca(2+)-activated K<sup>+</sup>-channel Aug 1992

19/6/33 (Item 33 from file: 155) 14645123 PMID: 12560073  
Molecular dynamics simulations of a K<sup>+</sup> channel blocker: Tc1 toxin from Thyus cambridgei. Jan 30 2003

19/6/34 (Item 34 from file: 155) 14194806 PMID: 11864985  
Mapping the binding site of a human ether-a-go-go-related gene-specific peptide toxin (ErgTx) to the channel's outer vestibule. May 10 2002

19/6/35 (Item 35 from file: 5) 0013745123 BIOSIS NO.: 200200338634  
Mapping the binding site of a human ether-a-go-go-related gene-specific peptide toxin (ErgTx) to the channel's outer vestibule 2002

19/6/36 (Item 36 from file: 155) 05208395 PMID: 1702643  
Mapping the receptor site for charybdotoxin, a pore-blocking potassium channel inhibitor. Dec 1990

19/6/37 (Item 37 from file: 155) 13436286 PMID: 10398697  
A marine snail neurotoxin shares with scorpion toxins a convergent mechanism of blockade on the pore of voltage-gated K channels. Jul 1999

19/6/38 (Item 38 from file: 5) 0012127174 BIOSIS NO.: 199900386834  
A marine snail neurotoxin shares with scorpion toxins a convergent mechanism of blockade on the pore of voltage-gated K channels 1999

19/6/39 (Item 39 from file: 155) 14633777 PMID: 12538890  
Motions and structural variability within toxins: implication for their use as scaffolds for protein engineering. Feb 2003

19/6/40 (Item 40 from file: 5) 0014168498 BIOSIS NO.: 200300125608  
Motions and structural variability within toxins: Implication for their use as scaffolds for protein engineering. 2003

19/6/41 (Item 41 from file: 155) 11833908 PMID: 9092804  
NMR solution structure of a two-disulfide derivative of charybdotoxin: 0 structural evidence for conservation of scorpion toxin alpha/beta motif and its hydrophobic side chain packing. Apr 1 1997

19/6/42 (Item 42 from file: 155) 11322494 PMID: 8645186  
A novel structural class of K<sup>+</sup>-channel blocking toxin from the scorpion Pandinus imperator. May 1 1996

19/6/43 (Item 43 from file: 5) 0012367417 BIOSIS NO.: 200000865730  
A point mutation in the maxi-K clone d5b forms a high affinity site for charybdotoxin 1999

19/6/44 (Item 44 from file: 155) 10274627 PMID: 8360176  
Purification, characterization, and biosynthesis of margatoxin, a component of Centruroides margaritatus venom that selectively inhibits voltage-dependent potassium channels. Sep 5 1993

19/6/45 (Item 45 from file: 5) 0008972783 BIOSIS NO.: 199336137199  
Purification, characterization, and biosynthesis of margatoxin, a component of Centruroides margaritatus venom that selectively inhibits voltage-dependent potassium channels 1993

19/6/46 (Item 46 from file: 155) 14164320 PMID: 11952787  
Purification, characterization and biosynthesis of paratubulin 3, a component of Parabuthus transvaalicus venom. Apr 2002

19/6/47 (Item 47 from file: 5) 0013693181 BIOSIS NO.: 200200286692  
Purification, characterization and biosynthesis of paratubulin 3, a component of Parabuthus transvaalicus venom 2002

19/6/48 (Item 48 from file: 155) 09024601 PMID: 1694175  
Purification and characterization of a unique, potent, peptidyl probe for the high conductance calcium-activated potassium channel from venom of the scorpion Buthus tamulus. Jul 5 1990

19/6/49 (Item 49 from file: 5) 0007279032 BIOSIS NO.: 199090063511  
PURIFICATION AND CHARACTERIZATION OF A UNIQUE POTENT PEPTIDYL PROBE FOR THE HIGH CONDUCTANCE CALCIUM-ACTIVATED POTASSIUM CHANNEL FROM VENOM OF THE SCORPION BUTHUS-TAMULUS 1990

19/6/50 (Item 50 from file: 155) 12065981 PMID: 9354615  
Purification, characterization, and synthesis of three novel toxins from the Chinese scorpion Buthus martensi, which act on K<sup>+</sup> channels. Nov 4 1997

19/6/51 (Item 51 from file: 155) 12568895 PMID: 9891977  
Purification and partial characterization of a 'short' insectotoxin-like peptide from the venom of the scorpion Parabuthus schlechteri. Dec 28 1998

19/6/52 (Item 52 from file: 5) 0011824566 BIOSIS NO.: 19990084226  
Purification and partial characterization of a 'short' insectotoxin-like peptide from the venom of the scorpion Parabuthus schlechteri 1998

19/6/53 (Item 53 from file: 155) 10018942 PMID: 1467342  
Progress in multidimensional NMR investigations of peptide and protein 3-D structures in solution. From structure to functional aspects. Sep-Oct 1992

19/6/54 (Item 54 from file: 5) 0008744162 BIOSIS NO.: 199395046428  
Progress in multidimensional NMR investigations of peptide and protein 3-D structures in solution: From structure to functional aspects 1992

19/6/55 (Item 55 from file: 155) 12054583 PMID: 9441593  
[The rational evolution of scorpion toxins] Ratsional'naya evolyutsiya toksinov iz yada skorpionov Sep 1997

19/6/56 (Item 56 from file: 5) 0011277181 BIOSIS NO.: 199800071428  
The rational evolution of scorpion toxins 1997

19/6/57 (Item 57 from file: 155) 12019132 PMID: 9306273  
Scorpion toxin block of the early K<sup>+</sup>-current (IKf) in rat dorsal root ganglion neurones. Sep 1 1997

19/6/58 (Item 58 from file: 5) 0011155973 BIOSIS NO.: 199799790033  
Scorpion toxin block of the early K<sup>+</sup>-current (IKf) in rat dorsal root ganglion neurones 1997

19/6/59 (Item 59 from file: 155) 15269022 PMID: 15049683  
Solution structure of BmBKTx1, a new BKCa1 channel blocker from the Chinese scorpion Buthus martensi Karsch. Apr 6 2004

19/6/60 (Item 60 from file: 5) 0014975010 BIOSIS NO.: 200400345799  
Solution structure of BmBKTx1, a new BKCa1 channel blocker from the Chinese scorpion *Buthus martensi* Karscht 2004

19/6/61 (Item 61 from file: 155) 14028726 PMID: 11790849  
Solution structure of a K(+) channel blocker from the scorpion *Tityus cambridgei*. Feb 2002

19/6/62 (Item 62 from file: 155) 11805876 PMID: 9062103  
Solution structure for Pandinus toxin K-alpha (PTX-K alpha), a selective blocker of A-type potassium channels. Mar 11 1997

19/6/63 (Item 63 from file: 155) 12075374 PMID: 9365990  
Solution structure of TsKapa, a charybotoxin-like scorpion toxin from *Tityus serrulatus* with high affinity for apamin-sensitive Ca(2+)-activated K+ channels. Nov 1997

19/6/64 (Item 64 from file: 5) 0010326748 BIOSIS NO.: 199598796581  
Solution structure of SHK toxin, a novel potassium channel inhibitor from a sea anemone 1996

19/6/65 (Item 65 from file: 155) 11968450 PMID: 9252467  
Sapadin B, a novel fly toxin, blocks macroscopic K+ currents in the GH3 rat pituitary cell line. Jul 1997

19/6/66 (Item 66 from file: 5) 001067804 BIOSIS NO.: 199799701864  
Sapadin B, a novel fly toxin, blocks macroscopic K+ currents in the GH-3 rat pituitary cell line 1997

19/6/67 (Item 67 from file: 5) 0012839066 BIOSIS NO.: 200100010905  
Structure determinants of scorpion toxin affinity: The charybotoxin (alpha-KTX) family of K+-channel blocking peptides BOOK TITLE: Reviews of Physiology Biochemistry and Pharmacology 2000

19/6/68 (Item 68 from file: 155) 12909571 PMID: 10857399  
Structural determinants of scorpion toxin affinity: the charybotoxin (alpha-KTX) family of K(+) channel blocking peptides. 2000

19/6/69 (Item 69 from file: 155) 12782762 PMID: 10707030  
Structural and functional differences of two toxins from the scorpion *Pandinus imperator*. Mar 1 2000

19/6/70 (Item 70 from file: 5) 0012447874 BIOSIS NO.: 2000000166187  
Structural and functional differences of two toxins from the scorpion *Pandinus imperator* 2000

19/6/71 (Item 71 from file: 5) 0010988956 BIOSIS NO.: 199799623016  
Structural analysis of a two disulfide bridge analogue of a scorpion toxin 1997

19/6/72 (Item 72 from file: 155) 10377674 PMID: 8253752  
Synthesis and characterization of kalitoxin. Is the 26-32 sequence essential for potassium channel recognition? Dec 15 1993

19/6/73 (Item 73 from file: 155) 10341798 PMID: 7693459  
Synthesis of charybotoxin and of two N-terminal truncated analogues. Structural and functional characterisation. Oct 1 1993

19/6/74 (Item 74 from file: 155) 11245554 PMID: 8547346  
Synthesis and structural characterisation of analogues of the potassium channel blocker charybotoxin. Jan 4 1996

19/6/75 (Item 75 from file: 155) 11180566 PMID: 7576659  
Topology of the pore-region of a K+ channel revealed by the NMR-derived structures of scorpion toxins. Nov 1995

19/6/76 (Item 76 from file: 155) 10582194 PMID: 7514038  
Tremorgenic indole alkaloids potently inhibit smooth muscle high-conductance calcium-activated potassium channels. May 17 1994

19/6/77 (Item 77 from file: 155) 14501863 PMID: 12445473  
Two novel toxins from the Amazonian scorpion *Tityus cambridgei* that block Kv1.3 and Shaker B K(+) channels with distinctly different affinities. Dec 16 2002

19/6/78 (Item 78 from file: 155) 12343405 PMID: 9655636  
Two similar peptides from the venom of the scorpion *Pandinus imperator*, one highly effective blocker and the other inactive on K+ channels. May 1998

19/6/79 (Item 79 from file: 5) 0011532158 BIOSIS NO.: 1998000326405  
Two similar peptides from the venom of the scorpion *Pandinus imperator*, one highly effective blocker and the other inactive on K+ channels 1998

19/6/80 (Item 80 from file: 155) 11603837 PMID: 8913348  
Three new toxins from the scorpion *Pandinus imperator* selectively block certain voltage-gated K+ channels. Nov 1996

19/6/81 (Item 81 from file: 5) 0010694277 BIOSIS NO.: 199799328337  
Three new toxins from the scorpion *Pandinus imperator* selectively block certain voltage-gated K+ channels 1996

24/6/1 (Item 1 from file: 155) 11928755 PMID: 9208943  
Anti-insect toxin 5 (AaIT5) from *Androctonus australis*. Jun 1 1997

24/6/2 (Item 2 from file: 5) 0010977935 BIOSIS NO.: 199799611995  
Anti-insect toxin 5 (AaIT5) from *Androctonus australis* 1997

24/6/3 (Item 3 from file: 357) 0206373 DBR Accession No.: 97-01494  
Control of insects, acari and nematodes - recombinant Autographa californica nuclear-polyhedrosis virus vector-mediated scorpion, wasp, snail, mite or spider venom toxin gene expression for use as a biological control agent 1996

24/6/4 (Item 4 from file: 155) 11146500 PMID: 8533143  
Positive cooperativity among insecticidal scorpion neurotoxins. Aug 1995

24/6/5 (Item 5 from file: 5) 0009989127 BIOSIS NO.: 199598456960  
Positive cooperativity among insecticidal scorpion neurotoxins 1995

24/6/6 (Item 6 from file: 155) 12339622 PMID: 9652392  
A depressant insect-selective toxin analog from the venom of the scorpion *Leiurus quinquestriatus hebraeus*-purification and structure/function characterization. May 15 1998

24/6/7 (Item 7 from file: 5) 0011517804 BIOSIS NO.: 199800312051  
A depressant insect-selective toxin analog from the venom of the scorpion *Leiurus quinquestriatus hebraeus*: Purification and structure/function characterization 1998

24/6/8 (Item 8 from file: 155) 10940776 PMID: 7722081  
Direct effects of recombinant nuclear polyhedrosis viruses on selected nontarget organisms. Apr 1995

24/6/9 (Item 9 from file: 5) 0009776510 BIOSIS NO.: 199598244443  
Direct effects of recombinant nuclear polyhedrosis viruses on selected nontarget organisms 1995

24/6/10 (Item 10 from file: 5) 0014296947 BIOSIS NO.: 200300255666  
Effect of signal sequence and promoter on the speed of action of a genetically modified Autographa californica nucleopolyhedrovirus expressing the scorpion toxin LqhIT2. 2003

24/6/11 (Item 11 from file: 357) 0313302 DBR Accession No.: 2003-14442  
Effect of signal sequence and promoter on the speed of action of a genetically modified Autographa californica nucleopolyhedrovirus expressing the scorpion toxin LqhIT2 - scorpion venom toxin expression in nuclear-polyhedrosis virus for biological control agent strain improvement 2003

24/6/12 (Item 12 from file: 155) 12084031 PMID: 9374595  
Interactions of recombinant and wild-type baculoviruses with classical insecticides and pyrethroid-resistant tobacco budworm (Lepidoptera: Noctuidae). Oct 1997

24/6/13 (Item 13 from file: 5) 0009921314 BIOSIS NO.: 199598389147  
Insect sodium channel as the target for insect-selective neurotoxins from scorpion venom BOOK TITLE: ACS Symposium Series; Molecular action of insecticides on ion channels 1995

24/6/14 (Item 14 from file: 5) 0009184411 BIOSIS NO.: 199497205696  
The insect sodium channel as the target for the insect selective neurotoxins from scorpion venom 1994

24/6/15 (Item 15 from file: 5) 0010929668 BIOSIS NO.: 199799563728  
Insect tolerance to a neurotoxic polypeptide: Pharmacokinetic and pharmacodynamic aspects 1997

24/6/16 (Item 16 from file: 155) 14493200 PMID: 11782289  
Isolation and characterization of a novel lepidopteran-selective toxin from the venom of South Indian red scorpion, *Mesobuthus tamulus*. 2001

24/6/17 (Item 17 from file: 5) 0013793695 BIOSIS NO.: 200200387406  
Isolation and characterization of a novel lepidopteran-selective toxin from the venom of South Indian red scorpion, *Mesobuthus tamulus* 2001

24/6/18 (Item 18 from file: 357) 0264969 DBR Accession No.: 2001-04723  
New polynucleotides encoding scorpion venom potassium channel-agonist proteins for producing e.g. of insect-tolerant transgenic plants for controlling insect pest damage and parasitic worm infections - scorpion venom potassium channel-agonist protein genes useful for constructing transgenic plant with insect resistance 2000

24/6/19 (Item 19 from file: 357) 0264374 DBR Accession No.: 2001-04128  
New isolated polynucleotide encoding a scorpion toxin for treating epilepsy, degenerative disorders such as Huntington's disease, and neuronal death following stroke, and for creating plants that are insect tolerant - transgenic plant construction with insect resistance and gene therapy 2000

24/6/20 (Item 20 from file: 5) 0008664579 BIOSIS NO.: 199345095567  
Potential of baculoviruses expressing a scorpion toxin and an esterase in agriculture 1993

24/6/21 (Item 21 from file: 357) 0153538 DBR Accession No.: 93-11590

Potential of baculo viruses expressing a scorpion toxin and an esterase in agriculture - use of recombinant baculovirus with a Heliothis virescens juvenile-hormone-esterase as an insect biological control agent (conference abstract) 1993

24/6/22 (Item 22 from file: 155) 12317587 PMID: 9627406  
Rapid purification and molecular modeling of AaIT peptides from venom of *Androctonus australis*. 1998

24/6/23 (Item 23 from file: 5) 0011525675 BIOSIS NO.: 199800319922  
Rapid purification and molecular modeling of AaIT peptides from venom of *Androctonus australis* 1998

24/6/24 (Item 24 from file: 5) 0014912774 BIOSIS NO.: 200400283531  
Scorpion toxins 2004

24/6/25 (Item 25 from file: 5) 0007297260 BIOSIS NO.: 199090081739  
A SCORPION VENOM NEUROTOXIN PARALYTIC TO INSECTS THAT AFFECTS SODIUM CURRENT INACTIVATION PURIFICATION PRIMARY STRUCTURE AND MODE OF ACTION 1990

24/6/26 (Item 26 from file: 155) 09075706 PMID: 2383565  
A scorpion venom neurotoxin paralytic to insects that affects sodium current inactivation: purification, primary structure, and mode of action. Jun 26 1990

24/6/27 (Item 27 from file: 5) 0007645125 BIOSIS NO.: 199191028016  
THE TOLERANCE OF LEPIDOPTEROUS LARVAE TO AN INSECT SELECTIVE NEUROTOXIN 1990

24/6/28 (Item 28 from file: 5) 0009069061 BIOSIS NO.: 199497090346  
Variability among insect sodium channels revealed by selective neurotoxins 1994

24/6/29 (Item 29 from file: 5) 0010988896 BIOSIS NO.: 199799623056  
Two novel short insectotoxins from the Asian scorpions *Buthus martensii* and *Buthus tamulus* 1997

24/7/6 (Item 6 from file: 155) DIALOG(R)File 155.MEDLINE(R)(c) format only 2005 Dialog. All rts. reserv.  
12339622 PMID: 9652392  
A depressant insect-selective toxin analog from the venom of the scorpion *Leiurus quinquestriatus hebraeus*-purification and structure/function characterization.  
Moskowitz H; Hermann R; Jones A D; Hammock B D  
Department of Entomology, University of California, Davis 95616, USA.  
European journal of biochemistry / FEBS (GERMANY) May 15 1998, 254 (1) p44-9, ISSN 0014-2956 Journal Code: 0107600 Contract/Grant No.: P 30 ES05707; ES; NIEHS Publishing Model Print Document type: Journal Article  
Languages: ENGLISH Main Citation Owner: NLM Record type: MEDLINE; Completed

The scorpion venom-derived excitatory and depressant insect-selective polypeptide neurotoxins modify sodium conductance in insect neuronal membranes and differ greatly in their primary structures and symptoms induced in blow fly larvae. We report here the purification and characterization of a new insect selective toxin, LqhIT5. LqhIT5 is more similar to the excitatory toxins in its mode of action and the depressant toxins in its primary structure. This toxin is a single polypeptide composed of 61 amino acids that are cross linked by four disulfide bonds. When LqhIT5 is injected into blow fly larvae, a fast contraction paralysis occurs without depressant activity. No mammalian toxicity was detected by subcutaneous or intracranial injections of this toxin into mice. Sequence comparison of LqhIT5 and known depressant toxins shows a high degree of similarity among the amino acids located on the C-terminus of the toxins. However, there are some clear differences in the amino acids located close to the N-terminus of the toxins. By the aid of homology modeling, we demonstrated that these amino acids have the same orientation in the tertiary structure of the molecule and are exposed to the environment. The change in the mode of action of LqhIT5 (no depressant activity) by substitutions of a few amino acids located on a specific exposed area of the toxin shed a new light on the structure/function relationship of scorpion toxins. These results caution that similarity in the mechanism of action of scorpion toxins does not always follow from an overall similarity in sequence.  
Record Date Created: 19980728 Record Date Completed: 19980728

26/6/1 (Item 1 from file: 155) 11928755 PMID: 9208943  
Anti-insect toxin 5 (AaIT5) from *Androctonus australis*. Jun 1 1997

26/6/2 (Item 2 from file: 5) 0010977935 BIOSIS NO.: 199799611995  
Anti-insect toxin 5 (AaIT5) from *Androctonus australis* 1997

26/6/3 (Item 3 from file: 155) 12339622 PMID: 9652392  
A depressant insect-selective toxin analog from the venom of the scorpion *Leiurus quinquestriatus hebraeus*-purification and structure/function characterization. May 15 1998

26/6/4 (Item 4 from file: 5) 0011517804 BIOSIS NO.: 199800312051  
A depressant insect-selective toxin analog from the venom of the scorpion *Leiurus quinquestriatus hebraeus*: Purification and structure/function characterization 1998

26/6/5 (Item 5 from file: 5) 0014296947 BIOSIS NO.: 200300255666  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolyhedrovirus expressing the scorpion toxin LqhIT2. 2003

26/6/6 (Item 6 from file: 357) 0313302 DBR Accession No.: 2003-14442  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolyhedrovirus expressing the scorpion toxin LqhIT2 - scorpion venom toxin expression in nuclear-polyhedrosis virus for biological control agent strain improvement 2003

26/6/7 (Item 7 from file: 5) 0007297260 BIOSIS NO.: 199090081739  
A SCORPION VENOM NEUROTOXIN PARALYTIC TO INSECTS THAT AFFECTS SODIUM CURRENT INACTIVATION PURIFICATION PRIMARY STRUCTURE AND MODE OF ACTION 1990

26/6/8 (Item 8 from file: 155) 09075706 PMID: 2383565  
A scorpion venom neurotoxin paralytic to insects that affects sodium current inactivation: purification, primary structure, and mode of action. Jun 26 1990

26/6/9 (Item 9 from file: 5) 0009069061 BIOSIS NO.: 199497090346  
Variability among insect sodium channels revealed by selective neurotoxins 1994

28/6/1 (Item 1 from file: 357) 0313302 DBR Accession No.: 2003-14442  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolyhedrovirus expressing the scorpion toxin LqhIT2 - scorpion venom toxin expression in nuclear-polyhedrosis virus for biological control agent strain improvement 2003

29/6/1 (Item 1 from file: 5) 0014419169 BIOSIS NO.: 200300377888  
Scorpion toxins 2003

29/6/2 (Item 1 from file: 357) 0313302 DBR Accession No.: 2003-14442  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolyhedrovirus expressing the scorpion toxin LqhIT2 - scorpion venom toxin expression in nuclear-polyhedrosis virus for biological control agent strain improvement 2003

29/6/3 (Item 2 from file: 357) 0256837 DBR Accession No.: 2000-11327  
New nucleic acid fragment encoding a scorpion toxin that is potassium channel-agonist, useful for creating transgenic plants that are more insect-tolerant - method is useful for producing transgenic plant with insect resistance 2000

33/6/1 (Item 1 from file: 5) 0014296947 BIOSIS NO.: 200300255666  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolyhedrovirus expressing the scorpion toxin LqhIT2. 2003

33/6/2 (Item 1 from file: 357) 0313302 DBR Accession No.: 2003-14442  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolyhedrovirus expressing the scorpion toxin LqhIT2 - scorpion venom toxin expression in nuclear-polyhedrosis virus for biological control agent strain improvement 2003

34/6/1 (Item 1 from file: 5) 0014419169 BIOSIS NO.: 200300377888  
Scorpion toxins 2003

34/6/2 (Item 2 from file: 5) 0014296947 BIOSIS NO.: 200300255666  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolyhedrovirus expressing the scorpion toxin LqhIT2. 2003

34/6/3 (Item 1 from file: 357) 0313302 DBR Accession No.: 2003-14442  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolyhedrovirus expressing the scorpion toxin LqhIT2 - scorpion venom toxin expression in nuclear-polyhedrosis virus for biological control agent strain improvement 2003

34/6/4 (Item 2 from file: 357) 0256837 DBR Accession No.: 2000-11327  
New nucleic acid fragment encoding a scorpion toxin that is potassium channel-agonist, useful for creating transgenic plants that are more insect-tolerant - method is useful for producing transgenic plant with insect resistance 2000

43/6/1 (Item 1 from file: 155) 09594018 PMID: 1796479  
The cDNA sequence of a depressant insect selective neurotoxin from the scorpion *Buthus judaicus*. 1991

43/6/2 (Item 2 from file: 5) 0008168878 BIOSIS NO.: 199293011769  
THE CDNA SEQUENCE OF A DEPRESSANT INSECT SELECTIVE NEUROTOXIN FROM THE SCORPION *BUTHOTUS-JUDAICUS* 1991

43/6/3 (Item 3 from file: 155) 12780322 PMID: 10708793  
Cloning and characterization of a cDNA sequence encoding the precursor of a chbrototoxin-like peptide from the Chinese scorpion *Buthus martensii* Karsch. Aug 2000

43/6/4 (Item 4 from file: 5) 0012519314 BIOSIS NO.: 2000000237627  
Cloning and characterization of a cDNA sequence encoding the precursor of a chbrototoxin-like peptide from the Chinese Scorpion *Buthus martensii* Karsch 2000

- 43/6/5 (Item 5 from file: 5) 0014778747 BIOSIS NO.: 200400145408  
Cytotoxic and apoptotic effects of scorpion *Leiurus quinquestriatus* venom on 293T and C2C12 eukaryotic cell lines. 2003
- 43/6/6 (Item 6 from file: 155) 10074904 PMID: 8431601  
Depressant insect selective neurotoxins from scorpion venom: chemistry, action, and gene cloning. 1993
- 43/6/7 (Item 7 from file: 5) 0011914925 BIOSIS NO.: 199900174585  
Dynamic diversification from a putative common ancestor of scorpion toxins affecting sodium, potassium, and chloride channels 1999
- 43/6/8 (Item 8 from file: 5) 0007735556 BIOSIS NO.: 199191118447  
DESIGN SYNTHESIS AND FUNCTIONAL EXPRESSION OF A GENE FOR CHARYBOTOXIN A PEPTIDE BLOCKER OF POTASSIUM ION CHANNELS 1991
- 43/6/9 (Item 9 from file: 155) 12095528 PMID: 9395089  
Influence of a NH<sub>2</sub>-terminal extension on the activity of KTX2, a K<sup>+</sup>-channel blocker purified from *Androctonus australis* scorpion venom. Nov 3 1997
- 43/6/10 (Item 10 from file: 5) 0014296947 BIOSIS NO.: 200300255666  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolydnavirus expressing the scorpion toxin LqhIT2. 2003
- 43/6/11 (Item 11 from file: 357) 0313302 DBR Accession No.: 2003-14442  
Effect of signal sequence and promoter on the speed of action of a genetically modified *Autographa californica* nucleopolydnavirus expressing the scorpion toxin LqhIT2 - scorpion venom toxin expression in nuclear-polyhedrosis virus for biological control agent strain improvement. 2003
- 43/6/12 (Item 12 from file: 155) 12401430 PMID: 9714546  
Evidence for a new class of scorpion toxins active against K<sup>+</sup>-channels. Jul 24 1998
- 43/6/13 (Item 13 from file: 155) 11211987 PMID: 7498537  
Functional expression of an alpha anti-insect scorpion neurotoxin in insect cells and lepidopterous larvae. Dec 4 1995
- 43/6/14 (Item 14 from file: 5) 0010170013 BIOSIS NO.: 199698637846  
Functional expression of an alpha anti-insect scorpion neurotoxin in insect cells and lepidopterous larvae 1995
- 43/6/15 (Item 15 from file: 357) 0191434 DBR Accession No.: 96-02205  
Functional expression of an alpha anti-insect scorpion neurotoxin in insect cells and lepidopterous larvae - *Autographa californica* nuclear-polyhedrosis virus vector allows insecticide expression in insect cell culture or larva 1995
- 43/6/16 (Item 16 from file: 5) 0012082499 BIOSIS NO.: 199900342159  
Genomic organization of three neurotoxins active on small conductance Ca<sub>2</sub>--activated potassium channels from the scorpion *Buthus maritansi* Karsch 1999
- 43/6/17 (Item 17 from file: 155) 18326469 PMID: 15966742  
Genetic polymorphism and expression of a highly potent scorpion depressant toxin enable refinement of the effects on insect Na channels and illuminate the key role of Asn-58. Jun 28 2005
- 43/6/18 (Item 18 from file: 5) 0015265895 BIOSIS NO.: 200500172631  
Improved plant protective efficacy of a baculovirus using an early promoter to drive insect-specific neurotoxin expression 2005
- 43/6/19 (Item 19 from file: 155) 15422272 PMID: 15133045  
Molecular basis of the high insecticidal potency of scorpion alpha-toxins. Jul 23 2004
- 43/6/20 (Item 20 from file: 5) 0008366559 BIOSIS NO.: 199294068400  
MOLECULAR ANALYSIS OF CDNA AND THE TRANSCRIPT ENCODING THE DEPRESSANT INSECT SELECTIVE NEUROTOXIN OF THE SCORPION *LEIURUS-QUINQUESTRIATUS-HEBRAEUS* 1992
- 43/6/21 (Item 21 from file: 5) 0012310167 BIOSIS NO.: 200000028480  
Mortality and feeding of mid-stadium larvae of *Helicoverpa zea* and *Heliothis virescens* fed a wild strain or a recombinant strain of *Baculovirus heliothis* expressing an insect-specific toxin of the scorpion *Leiurus quinquestriatus hebraeus* 1999
- 43/6/22 (Item 22 from file: 155) 09709895 PMID: 1801321  
Nucleotide sequence and structure analysis of a cDNA encoding an alpha insect toxin from the scorpion *Leiurus quinquestriatus hebraeus*. 1991
- 43/6/23 (Item 23 from file: 5) 0008192856 BIOSIS NO.: 199293035747  
NUCLEOTIDE SEQUENCE AND STRUCTURE ANALYSIS OF A CDNA ENCODING AN ALPHA INSECT TOXIN FROM THE SCORPION *LEIURUS-QUINQUESTRIATUS-HEBRAEUS* 1991
- 43/6/24 (Item 24 from file: 155) 14817292 PMID: 12787033  
An 'Old World' scorpion beta-toxin that recognizes both insect and mammalian sodium channels. Jun 2003
- 43/6/25 (Item 25 from file: 5) 0013196447 BIOSIS NO.: 200100368286  
Persistence and distribution of wild-type and recombinant nucleopolydnaviruses in soil 2001
- 43/6/26 (Item 26 from file: 5) 0013009537 BIOSIS NO.: 200100181376  
Recombinant baculovirus insecticides 2000
- 43/6/27 (Item 27 from file: 357) 0108634 DBR Accession No.: 90-11325  
Recombinant DNA coding for insecticide toxin - including new scorpion venom toxin or spider venom toxin, and transgenic plant containing such DNA 1990
- 43/6/28 (Item 28 from file: 357) 0148040 DBR Accession No.: 93-06092  
Rapid isolation of full length cDNA clones by 'inverse PCR': purification of a scorpion cDNA family encoding alpha-neurotoxins - cDNA clone isolation by the inverse polymerase chain reaction 1993
- 43/6/29 (Item 29 from file: 5) 0012648094 BIOSIS NO.: 2000000366407  
Scorpion neurotoxins: Structure/function relationships and application in agriculture 2000
- 43/6/30 (Item 30 from file: 5) 0014419169 BIOSIS NO.: 200300377888  
Scorpion toxins 2003
- 43/6/31 (Item 31 from file: 5) 0012786805 BIOSIS NO.: 2000000505118  
Yields of occlusion bodies from *Heliothis virescens* (Lepidoptera: Noctuidae) and *Helicoverpa zea* (Lepidoptera: Noctuidae) larvae fed wild or recombinant strains of baculoviruses 2000
- 43/6/32 (Item 32 from file: 155) 14611132 PMID: 12467668  
Three polymorphic genes encoding a depressant toxin from the Egyptian scorpion *Leiurus quinquestriatus quinquestriatus*. Jan 2003
- 43/6/33 (Item 33 from file: 5) 0014463392 BIOSIS NO.: 200300105111  
Three polymorphic genes encoding a depressant toxin from the Egyptian scorpion *Leiurus quinquestriatus quinquestriatus*. 2003
- 48/6/1 (Item 1 from file: 5) 0014419169 BIOSIS NO.: 200300377888  
Scorpion toxins 2003
- 48/6/2 (Item 1 from file: 357) 0264969 DBR Accession No.: 2001-04723  
New polynucleotides encoding scorpion venom potassium channel-agonist proteins for producing e.g. of insect-tolerant transgenic plants for controlling insect pest damage and parasitic worm infections - scorpion venom potassium channel-agonist protein genes useful for constructing transgenic plant with insect resistance 2000
- 48/6/3 (Item 2 from file: 357) 0264374 DBR Accession No.: 2001-04128  
New isolated polynucleotide encoding a scorpion toxin for treating epilepsy, degenerative disorders such as Huntington's disease, and neuronal death following stroke, and for creating plants that are insect tolerant - transgenic plant construction with insect resistance and gene therapy 2000
- 48/7/2 (Item 1 from file: 357) DIALOG(R)File 357:Derwent Biotech Res. (c) 2005 Thomson Derwent & ISI. All rts. reserv.  
0264969 DBR Accession No.: 2001-04723 PATENT  
New polynucleotides encoding scorpion venom potassium channel-agonist proteins for producing e.g. of insect-tolerant transgenic plants for controlling insect pest damage and parasitic worm infections - scorpion venom potassium channel-agonist protein genes useful for constructing transgenic plant with insect resistance  
AUTHOR: Herrmann R; Lee J M; Wong J F  
CORPORATE SOURCE: Wilmington, DE, USA  
PATENT ASSIGNEE: Du-Pont 2000  
PATENT NUMBER: WO 2007/8958 PATENT DATE: 20001228 WPI ACCESSION NO.: 2001-071394 (2008)  
PRIORITY APPLIC. NO.: US 140227 APPLIC. DATE: 19990622  
NATIONAL APPLIC. NO.: WO 2000US17049 APPLIC. DATE: 20000621 LANGUAGE: English  
ABSTRACT: An isolated polynucleotide (I) (e.g. DNA or RNA) is claimed. (I) contains a nucleotide sequence selected from a nucleotide sequence (III) of at least 81 nucleotides selected from 10 sequences of 171-213 (N1)-(N10), a nucleotide sequence (III) encoding a protein of at least 27 amino acids selected from 10 sequences of 56-70 amino acids (P1)-(P10), or a complement of (II) or (IV). Also claimed are: a chimeric gene or vector (II); a host cell (yeast, bacterium, plant) containing (I) or (II); a virus containing (I); a protein of at least 27 amino acids; a method of obtaining a nucleic acid fragment encoding a K-channel agonist; a recombinant baculo virus expression vector; and a method for testing the activity of a K-channel agonist against insects. (I) is useful for creating transgenic plants which express K-channel modifiers, useful as a means for controlling insect pest by producing insect tolerance. In the prevention or treatment of insect pest damage and parasitic worm infections in animals and humans, the invention may also find use in creating specific new pesticides and anthelmintic drugs that are also non-toxic to humans, pets and livestock. (50pp)